

檔 號：
保存年限：

衛生福利部食品藥物管理署 函

地址：11561 臺北市南港區昆陽街161-2
號

聯絡人：黃寶萱

聯絡電話：02-27877670

傳真：02-3322-9527

電子信箱：m01156@fda.gov.tw

受文者：中華民國藥師公會全國聯合會

發文日期：中華民國110年5月17日

發文字號：FDA藥字第1101404774號

速別：普通件

密等及解密條件或保密期限：

附件：

主旨：國際醫藥法規協和會（ICH）S1B(R1)指引草案進入公開諮詢階段，請協助轉知所屬，彙整該指引有關建議並請於110年8月31日前惠復，請查照。

說明：

- 一、S1B(R1)工作係藥品之齧齒動物致癌性檢測（Addendum to the Guideline on Testing for Carcinogenicity of Pharmaceuticals）指引之修訂，該指引草案現進入法規諮詢階段公開徵求各界意見，請協助轉知所屬，並惠予彙整所屬建議，依意見彙整表提供中英文建議惠復。
- 二、S1B(R1)指引草案及意見彙整表請至本署ICH草案公開諮詢專區下載，路徑：首頁>業務專區>藥品>ICH專區>草案公開諮詢專區。草案相關資料亦可參考ICH Public Consultations (<https://www.ich.org/page/public-consultations>)。

正本：台灣藥學會、中華民國學名藥協會、中華民國西藥商業同業公會全國聯合會、中華民國藥劑生公會全國聯合會、台灣年輕藥師協會、台灣醫藥品法規學會、台灣研發型生技新藥發展協會、臺灣製藥工業同業公會、台灣藥品行銷暨管理協會、

台北市西藥商業同業公會、中華民國西藥代理商業同業公會、台北市西藥代理商
業同業公會、中華民國製藥發展協會、中華民國開發性製藥研究協會、中華民國
藥師公會全國聯合會、社團法人臺灣臨床藥學會、台北市進出口商業同業公會
副本：財團法人醫藥品查驗中心

電 2021/05/18 文
交 09:04 換 章

裝

線

INTERNATIONAL COUNCIL FOR HARMONISATION OF TECHNICAL
REQUIREMENTS FOR PHARMACEUTICALS FOR HUMAN USE

ICH HARMONISED GUIDELINE

**ADDENDUM TO THE GUIDELINE ON TESTING
FOR CARCINOGENICITY OF
PHARMACEUTICALS
S1B(R1)**

Draft version

Endorsed on 10 May 2021

Currently under public consultation

At Step 2 of the ICH Process, a consensus draft text or guideline, agreed by the appropriate ICH Expert Working Group, is transmitted by the ICH Assembly to the regulatory authorities of the ICH regions for internal and external consultation, according to national or regional procedures.

S1B(R1)
Document History

Code	History	Date
S1B(R1)*	Endorsement by the Members of the ICH Assembly under <i>Step 2</i> and release for public consultation.	10 May 2021

**This addendum is complementary to the S1 Guidelines (S1A, S1B and S1C(R2)) and is not intended to replace the existing S1B Guideline. At Step 4 of the ICH process, this addendum will be integrated with the S1B Guideline.*

Legal notice: *This document is protected by copyright and may, with the exception of the ICH logo, be used, reproduced, incorporated into other works, adapted, modified, translated or distributed under a public license provided that ICH's copyright in the document is acknowledged at all times. In case of any adaption, modification or translation of the document, reasonable steps must be taken to clearly label, demarcate or otherwise identify that changes were made to or based on the original document. Any impression that the adaption, modification or translation of the original document is endorsed or sponsored by the ICH must be avoided.*

The document is provided "as is" without warranty of any kind. In no event shall the ICH or the authors of the original document be liable for any claim, damages or other liability arising from the use of the document.

The above-mentioned permissions do not apply to content supplied by third parties. Therefore, for documents where the copyright vests in a third party, permission for reproduction must be obtained from this copyright holder.

ICH HARMONISED GUIDELINE
ADDENDUM TO THE GUIDELINE ON TESTING FOR
CARCINOGENICITY OF PHARMACEUTICALS

ICH S1B(R1)

ICH Consensus Guideline

TABLE OF CONTENTS

PREAMBLE	1
1. INTRODUCTION.....	1
1.1 Scope of the Addendum	1
1.2 Purpose of the Addendum	1
1.3 Background	1
2. A WEIGHT OF EVIDENCE APPROACH TO ASSESS THE HUMAN CARCINOGENIC	
POTENTIAL OF SMALL MOLECULE PHARMACEUTICALS	2
2.1 Factors to Consider for a WoE Assessment	3
2.2 Integration of WoE Factors for Assessing Human Carcinogenic Risk	4
2.3 Mouse Carcinogenicity Studies	4
3. CLARIFICATION OF CRITERIA FOR SELECTION OF HIGH DOSE FOR RASH2-TG	
MOUSE CARCINOGENICITY STUDIES.....	5
REFERENCES.....	5
APPENDIX 1: CASE STUDIES APPLYING THE WEIGHT OF EVIDENCE APPROACH..	7

1 **PREAMBLE**

2 This Addendum is to be used in close conjunction with ICH *S1A Guideline on the Need for*
3 *Carcinogenicity Studies for Pharmaceuticals, S1B Testing for Carcinogenicity of*
4 *Pharmaceuticals, and S1C(R2) Dose Selection for Carcinogenicity Studies*. The Addendum is
5 complementary to the S1 Guidelines.

6 **1. INTRODUCTION**

7 **1.1 Scope of the Addendum**

8 This Addendum covers all small molecule pharmaceuticals where carcinogenicity evaluations
9 are recommended as described in S1A.

10 **1.2 Purpose of the Addendum**

11 This Addendum expands the testing scheme for assessing human carcinogenic risk of small
12 molecule pharmaceuticals by introducing an additional approach that is not described in the
13 original S1B Guideline. This is an integrative approach that provides specific weight of
14 evidence [WoE] criteria that inform whether or not a 2-year rat study adds value in completing
15 a human carcinogenicity risk assessment. The Addendum also adds a plasma exposure ratio-
16 based approach for setting the high dose in the rasH2-Tg mouse model,¹ while all other aspects
17 of the recommendations for high dose selection in S1C(R2) Guideline would still apply.

18 Application of this integrative approach would reduce the use of animals in accordance with the
19 3Rs (reduce/refine/replace) principles, and shift resources to focus onto generating more
20 scientific mechanism-based carcinogenicity assessments, while promoting safe and ethical
21 development of new small molecule pharmaceuticals.

22 **1.3 Background**

23 While the S1B Guideline calls for flexibility in considering approaches to address
24 pharmaceutical carcinogenicity testing, the basic scheme generally recommends a long-term
25 rodent study which, in practice, is usually a 2-year study in rats, along with a second rodent
26 carcinogenicity study in mice (2-year or short-term study). Since publication of the ICH S1B
27 Guideline, scientific advances toward elucidation of mechanisms of tumorigenic action, greater
28 understanding of the limitations of rodent models, and several retrospective analyses of
29 pharmaceutical datasets indicate that 2-year rat carcinogenicity studies might not add value to
30 human carcinogenicity risk assessment in some cases and the carcinogenic potential could have

¹ The rasH2-Tg mouse was developed in the laboratory of Tatsuji Nomura of the Central Institute for Experimental Animals (1). The model is referred to in the S1B Guideline as the TgHras2 transgenic mouse. The official nomenclature for the model is CByB6F1-Tg(HRAS)2Jic which is maintained by intercrossing C57BL/6JJic-Tg(HRAS)2Jic hemizygous male mice with BALB/cByJJic female mice. The littermates derived from these intercrosses are the transgenic rasH2-Tg animals with the tg/wt genotype, and the wild type rasH2-Wt animals with a wt/wt genotype.

Since other short-term models mentioned in S1B have not gained significant use compared to rasH2-Tg over the past 20 years, pharmaceutical development experience with these models is far more limited. Therefore, other short-term carcinogenicity models referred to in S1B would not qualify for a plasma exposure ratio-based high dose selection.

It is appropriate to use wild-type rasH2-Wt littermates of rasH2-Tg mice for dose range-finding studies and for generating exposure data.

31 been assessed adequately based on a comprehensive assessment of all available
32 pharmacological, biological, and toxicological data (2-9).

33 To determine whether the conclusions from these retrospective analyses could be confirmed in
34 a real- world setting (i.e., prior to knowledge of the 2-year rat carcinogenicity study outcomes),
35 an independent international prospective study was conducted under ICH *S1(R1) RND*
36 *Proposed Change to Rodent Carcinogenicity Testing of Pharmaceuticals – Regulatory Notice*
37 *Document*. The conclusion from this prospective evaluation confirmed that an integrated WoE
38 approach could be used to adequately assess the human carcinogenic risk for certain
39 pharmaceuticals in lieu of conducting a 2-year rat study.²

40 In addition, an exposure ratio endpoint (based on animal to human plasma AUC) for high dose
41 selection in 2-year rodent studies as per ICH S1C(R2) has not been globally accepted for use
42 in the rasH2-Tg mouse study. Therefore, a comprehensive analysis was conducted to assess
43 exposures and outcomes in rasH2-Tg studies from available information.³ As described in
44 Section 3, the results of this analysis indicate that there is no value in exceeding a 50-fold
45 exposure ratio for high dose selection in this model.

46 **2. A WEIGHT OF EVIDENCE APPROACH TO ASSESS THE HUMAN** 47 **CARCINOGENIC POTENTIAL OF SMALL MOLECULE PHARMACEUTICALS**

48 Over the course of drug development, it is important for sponsors to develop a scientifically
49 robust strategy for carcinogenicity assessment that considers key biologic, pharmacologic, and
50 toxicologic information. The integrative WoE assessment approach described in sections 2.1
51 and 2.2 may support a conclusion that the test compound is either:

- 52 • likely to be carcinogenic in humans such that the product would be labeled accordingly
53 and any 2-year rat carcinogenicity studies would not add value; or
- 54 • likely not to be carcinogenic in humans such that a 2-year rat study would not add value
55 (may also not be carcinogenic in rats, or may likely be carcinogenic in rats but through

² Conduct and results of the prospective study will be summarized; ICH Website of RND and PEP updates will be cited; and future DRA manuscript pointed to. These new citations will appear in the Step 4 Version and this footnote modified.

³ The approach taken for determining an adequate exposure margin for high dose selection for the rasH2-Tg short-term model is similar to that described previously for the 2-year rat and mouse studies (10,11) and Hisada S, Tsubota K, et al (Manuscript in preparation) Survey of Available Data to Assess Tumorigenic Sensitivity of rasH2-Tg Mice and 2-year Rodent Models. Draft Summary: Results were analyzed from studies conducted for 50 drugs in the 6-month rasH2-Tg model and the 2-year rat, 15 of which were also evaluated in the 2-year mouse. For 13 studies concluded to be positive in rasH2-Tg, 6 genotoxic carcinogens were positive within 0.1 - 3-fold of the AUC exposure ratio or body surface area adjusted dose ratio (rodent:human), and 7 nongenotoxic carcinogens were positive all within 1 - 50-fold. Among those 7, three tested positive only at exposures evaluated that exceeded 25-fold. The rasH2-Tg model was 20-fold more sensitive to 10-fold less sensitive than the 2-yr rat or mouse among these 13 drugs that were tested in all 3 models, while 3 of the 13 drugs tested negative in the 2-year rat study. Eight of 37 drugs that tested negative in rasH2-Tg were evaluated at greater than 50-fold exposure ratios (60 to >200-fold). For 11 compounds testing positive in 2-year rat studies at exposure ratios of <25-fold, and testing negative in rasH2-Tg, high dose selection in rasH2-Tg was limited by maximum tolerated dose (MTD) at exposure ratios of <50-fold for 9 drugs, and for the other 2 drugs, exposure margins exceeded 50-fold. Human relevance of the tumorigenic potential observed in rats for these 11 drugs has been questioned. In conclusion, when high exposures are tolerated in rasH2-Tg mice, there appears to be some value in exceeding 25-fold, but the overall evidence indicates no benefit to exceeding a 50-fold exposure margin. (Note: this summary paragraph may be deleted upon publication of Hisada et al).

- 56 well recognized mechanisms known to be human irrelevant); or
57 • uncertain with respect to the carcinogenic potential for humans, and a 2-year rat
58 carcinogenicity study is likely to add value to human risk assessment.

59

60 In cases where the WoE assessment leads to a conclusion of uncertainty regarding human
61 carcinogenicity potential, the approach described in S1B of conducting a 2-year rat
62 carcinogenicity study together with a carcinogenicity assessment in mice (short term or 2-year
63 study) remains the most appropriate strategy.

64 **2.1 Factors to consider for a WoE assessment**

65 A WoE approach is based on a comprehensive assessment of the totality of data relevant to
66 carcinogenic potential available from public sources and from conventional drug development
67 studies. These factors include:

- 68 1) data that inform carcinogenic potential based on drug target biology and the primary
69 pharmacologic mechanism of the parent compound and active major human
70 metabolites. This includes drug target distribution in rat and human; available
71 information from genetically engineered models; human genetic association studies;
72 cancer gene databases; and carcinogenicity information available on the drug class,
73 2) results from secondary pharmacology screens for the parent compound and major
74 metabolites that inform off-target potential, especially those that inform carcinogenic
75 risk (e.g., binding to nuclear receptors),
76 3) histopathology data from repeated-dose toxicity studies completed with the test agent,
77 with particular emphasis on the long term rat study, including exposure margin
78 assessments of parent drug and major metabolites,⁴
79 4) evidence for hormonal perturbation, including knowledge of drug target and
80 compensatory endocrine response mechanisms; weight, gross and microscopic changes
81 in endocrine and reproductive organs from repeated-dose toxicity studies; and results
82 from reproductive toxicology studies,⁵
83 5) genetic toxicology study data using criteria from ICH S2(R1) Genotoxicity Testing and
84 Data Interpretation for Pharmaceuticals Intended for Human Use; equivocal
85 genotoxicity increases uncertainty with respect to the carcinogenic potential,
86 6) evidence of immune modulation in accordance with ICH S8 Immunotoxicity Studies
87 for Human Pharmaceuticals; it is generally recognized (12,13) that standard rat and
88 mouse carcinogenicity studies are not reliable for identifying this specific human risk.

⁴ Histopathology findings from long term rat toxicity studies of particular interest for identifying carcinogenic potential in a 2-year rat study include cellular hypertrophy, cellular hyperplasia, persistent tissue injury and/or chronic inflammation, foci of cellular alteration, preneoplastic changes, and tumors. It is important to provide an understanding of the likely pathogenesis, and/or address the human relevance of such findings. While long term rat toxicity study data are shown to be of highest value for assessing the likely outcome and value of conducting a 2-year rat study, short term rat studies can sometimes also provide histopathologic conclusions of value.

Data from long term toxicity studies in non-rodents and mice may also be useful for providing additional context on the human relevance of rat study findings (e.g., species-specific mechanistic differences) and whether there is value in conducting a 2-yr rat study.

⁵ If microscopic changes in endocrine and reproductive tissues including atrophy, hypertrophy, hyperplasia are observed, or statistically and biologically significant test article associated endocrine or reproductive organ weight changes are observed this may be considered evidence of functional hormonal perturbation even when changes in hormone levels are not documented. Such findings may be suggestive of potential carcinogenic risk unless investigated for human relevance and demonstrated otherwise.

89 The above WoE factors may be sufficient to conclude whether or not a 2-year rat study would
 90 add value. However, where one or more WoE factors may be inconclusive or indicate a
 91 concern for carcinogenicity, the Sponsor can conduct investigations that could inform human
 92 relevance of the potential risk. Possible approaches may include, but are not limited to:

- 93 1) additional investigational studies, or analyses of specimens collected from prior studies
 94 (e.g., special histochemical stains, molecular biomarkers, serum hormone levels,
 95 further characterization of immunomodulation, alternative *in vitro* or *in vivo* test
 96 systems, data from emerging technologies, etc.), and
- 97 2) clinical data generated to inform human mechanistic relevance at therapeutic doses and
 98 exposures (e.g., urine drug concentrations and evidence of crystal formation; targeted
 99 measurements of clinical plasma hormonal alterations; human imaging data, etc.).

100 2.2 Integration of WoE Factors for Assessing Human Carcinogenic Risk

101 An integrated analysis of the WoE factors described above determines whether or not a
 102 standard 2-year rat study would contribute to the human carcinogenic risk assessment. While
 103 all factors will contribute to the integrated analysis, the relative importance of each factor will
 104 vary depending on the specific molecule being considered. A summary of key outcomes and
 105 examples based on the experience accrued during the ICH S1 RND study (*S1(R1) RND*
 106 *Proposed Change to Rodent Carcinogenicity Testing of Pharmaceuticals – Regulatory Notice*
 107 *Document*), are provided in Appendix 1 demonstrating how the WoE factors could be
 108 integrated in determining the need for a 2-year rat study.

109 Experience from the ICH S1 RND study indicates that an established profile of other
 110 compound(s) in a drug class contributes substantially to assessing human carcinogenic risk
 111 associated with modulation of the pharmacologic target. Compounds with novel drug targets
 112 (i.e., first-in-class) are, nevertheless, considered eligible for an integrative WoE-based
 113 approach. For such candidates, a higher evidentiary standard is expected to establish that there
 114 is no cause-for-concern in regard to target biology. Appendix 1 provides an example where a
 115 WoE assessment led to a conclusion that a 2-year rat study would not add value to human
 116 carcinogenic risk assessment for a drug inhibiting a novel target.

117 When the WoE assessment concludes that conduct of a 2-year rat study is not warranted, the
 118 Sponsor should seek alignment with the Drug Regulatory Agency [DRA] of each region where
 119 marketing approval is sought. When a sponsor decides to conduct a 2-year rat study in
 120 accordance with ICH S1B, there is no obligation to seek concurrence nor to document their
 121 rationale with each DRA.

122 2.3 Mouse Carcinogenicity Studies

123 A carcinogenicity study in mice, either 2-year or a short-term transgenic model as specified in
 124 ICH S1B, remains a recommended component of a carcinogenicity assessment plan, even for
 125 those compounds where the integrated WoE assessment indicates a 2-year rat study would not
 126 contribute significant value.⁶ However, in some cases, for example, when the WoE evaluation

⁶ The WoE approach described for the rat is not appropriate for eliminating the mouse as a second rodent carcinogenicity species because: (1) 6-month chronic toxicity studies are not generally conducted with mice so the WoE approach cannot be implemented and no database is available to confirm this approach, (2) the results of carcinogenicity studies in mice will often provide different outcomes from the corresponding rat carcinogenicity study, so a direct extrapolation cannot be made, and (3) a 6-month rasH2-Tg mouse has been adopted as an acceptable carcinogenicity study model.

When the WoE evaluation indicates the 2-year rat study adds no value, a carcinogenicity study in mice (either 2-year or short-term) is also not recommended in the EU.

127 strongly indicates no carcinogenic risk to humans and data indicate that only subtherapeutic,
 128 pharmacologically inactive drug exposures can be achieved in the mouse, it may not be
 129 appropriate to conduct any mouse carcinogenicity study.

130 **3. CLARIFICATION ON CRITERIA FOR SELECTION OF THE HIGH DOSE FOR**
 131 **RASH2-TG MOUSE CARCINOGENICITY STUDIES**

132 In practice, a plasma exposure (AUC) ratio for high dose selection in the absence of dose
 133 limiting toxicity or appropriate use of other dose setting criteria as outlined in ICH S1C(R2) in
 134 this model, has not been globally accepted as an endpoint. Therefore, available data from
 135 experience with 50 compounds evaluated in the rasH2-Tg mouse model were analyzed and the
 136 conclusion reached that there was no value in exceeding a 50-fold plasma AUC exposure ratio
 137 (rodent:human) to support carcinogenicity assessment. Therefore, all criteria for selection of
 138 the high dose for carcinogenicity studies as specified in S1C(R2) for 2-year rodent studies are
 139 applicable to rasH2-Tg, including an AUC plasma exposure ratio, except that the exposure
 140 ratio will be 50-fold in rasH2-Tg rather than 25-fold as for 2-year studies conducted in wild
 141 type rodents. All other aspects of S1C(R2) remain applicable to rasH2-Tg.

142 **REFERENCES**

- 143 (1) Saitoh A, Kimura M, Takahashi R, Yokoyama M, Nomura T, Izawa M et al. Most
 144 tumors in transgenic mice with human c-Ha-ras gene contained somatically activated
 145 transgenes. *Oncogene* 1990;5(8):1195-200.
- 146 (2) Van Oosterhout JPJ, Van der Laan JW, De Waal EJ, Olejniczak K, Hilgenfeld M,
 147 Schmidt V et al. The utility of two rodent species in carcinogenic risk assessment of
 148 pharmaceuticals in Europe. *Reg Toxicol Pharmacol* 1997;25:6-17.
- 149 (3) Contrera JF, Jacobs AC, DeGeorge JJ. Carcinogenicity testing and the evaluation of
 150 regulatory requirements for pharmaceuticals. *Reg Toxicol Pharmacol* 1997;25:130-45.
- 151 (4) Reddy MV, Sistare FD, Christensen JS, DeLuca JG, Wollenberg GK, DeGeorge JJ. An
 152 evaluation of chronic 6- and 12-month rat toxicology studies as predictors of 2-year
 153 tumor outcome. *Vet Pathol* 2010;47:614-29.
- 154 (5) Sistare FD, Morton D, Alden C, Christensen J, Keller D, De Jonghe S et al. An analysis
 155 of pharmaceutical experience with decades of rat carcinogenicity testing: support for a
 156 proposal to modify current regulatory guidelines. *Toxicol Pathol* 2011;39:716-44.
- 157 (6) Alden CL, Lynn A, Bourdeau A, Morton D, Sistare FD, Kadambi VJ et al. A critical
 158 review of the effectiveness of rodent pharmaceutical carcinogenesis testing in
 159 predicting for human risk. *Vet Pathol* 2011;48:772-84.
- 160 (7) Friedrich A, Olejniczak K. Evaluation of carcinogenicity studies of medicinal products
 161 for human use authorised via the European centralised procedure (1995-2009). *Reg*
 162 *Toxicol Pharmacol* 2011;60:225-48.
- 163 (8) Van der Laan JW, Kasper P, Lima BS, Jones DR, Pasanen M. Critical analysis of
 164 carcinogenicity study outcomes. Relationship with pharmacological properties. *Crit*
 165 *Rev Toxicol* 2016;46:587-614.
- 166 (9) Van der Laan JW, Buitenhuis WHW, Wagenaar L, Soffers AEMF, Van Someren EP,
 167 Krul CAM et al. Prediction of the carcinogenic potential of human pharmaceuticals
 168 using repeated dose toxicity data and their pharmacological properties. *Frontiers in*
 169 *Medicine* 2016;3:45. doi: 10.3389/fmed2016.00045

ICH S1B(R1) Guideline

- 170 (10) Contrera JF, Jacobs AC, Prasanna HR, Mehta M, Schmidt WJ, De George J. A
171 systemic exposure-based alternative to the maximum tolerated dose for carcinogenicity
172 studies of human therapeutics. *J Am Coll Toxicol* 1995;14:1–10.
- 173 (11) Bergman K, Olofsson IM, Sjöberg P. Dose selection for carcinogenicity studies of
174 pharmaceuticals: systemic exposure to phenacetin at carcinogenic dosage in the rat. *Reg
175 Toxicol Pharmacol* 1998;28:226–29.
- 176 (12) Bugelski PJ, Volk A, Walker MA, Krayner JH, Martin P, Descotes J. Critical review
177 of preclinical approaches to evaluate the potential of immunosuppressive drugs to
178 influence human neoplasia. *Int J Toxicol* 2010;29:435-66.
- 179 (13) Lebrech H, Brennan FR, Haggerty H, Herzyk D, Kamperschroer C, Maier CC et al.
180 HESI/FDA workshop on immunomodulators and cancer risk assessment: Building
181 blocks for a weight-of-evidence approach. *Reg Toxicol Pharmacol* 2016;75: 72-80.

182 **APPENDIX 1: CASE STUDIES APPLYING THE WEIGHT OF EVIDENCE**
 183 **APPROACH**

184
 185 **Preamble**
 186

187 One outcome of the ICH S1 RND study was the recognition that programs with the following
 188 WoE attributes are more likely to support a conclusion that the results of a 2-year rat study
 189 would not contribute value to human carcinogenicity risk assessment.

- 190 • Target biology is well characterized and not associated with cellular pathways known
 191 to be involved with human cancer development. Often, the pharmaceutical target was
 192 non-mammalian and carcinogenicity data were available with the pharmacologic drug
 193 class.
- 194 • Results from chronic toxicity studies indicate no hyperplastic, hypertrophic, atypical
 195 cellular alterations, or degenerative/regenerative changes noted without adequate
 196 explanation of pathogenesis or human relevance, indicative of no on- or off-target
 197 potential of carcinogenic concern;
- 198 • No perturbation of endocrine and reproductive organs observed, or endocrine findings
 199 adequately explained with respect to potential human relevance;
- 200 • No identified concerns from secondary pharmacology screens intended to inform off-
 201 target potential for the pharmaceutical
- 202 • No evidence of immune modulation or immunotoxicity based on target biology and
 203 repeat dose toxicology studies
- 204 • The overall assessment of genotoxic potential is concluded to be negative based on
 205 criteria from ICH S2(R1) Guidance.

206
 207 Although rasH2-Tg mouse study results were recommended when available as a WoE element
 208 in the initial RND, they did not significantly contribute to the prediction of the 2-year rat
 209 carcinogenicity study outcome. Therefore, a rasH2-Tg mouse study is not expected to be
 210 completed to support a WoE assessment. However, if rasH2-Tg mouse study results are
 211 available, they should be discussed in the assessment.

212
 213 A series of case studies are provided to illustrate the application of the WoE approach. These
 214 cases are provided for illustrative purposes only and are not intended as guidance to indicate
 215 the sufficiency of data to support a WoE assessment. Cases 1 and 2 describe the key WoE
 216 factors for that pharmaceutical and how the data were integrated to conclude that a 2-year rat
 217 study would not add value to the assessment of carcinogenic risk. In contrast to these cases,
 218 Case 3 describes how data from the WoE factors were integrated to conclude that the
 219 carcinogenic potential for humans was uncertain, and a 2-year rat carcinogenicity study was
 220 likely to add value to human risk assessment. Case 4 describes a molecule for which a 2-year
 221 rat carcinogenicity study was concluded to not contribute value to human carcinogenicity
 222 assessment despite there being no data available for other molecules within the pharmacologic
 223 class.

224
 225 **Case 1: A small molecule inhibitor against a non-mammalian target**
 226

227 **Prospective WoE Assessment: Concluded by all DRAs and Sponsor as likely not to be carcinogenic**
 228 **in both rats or humans such that a 2-year rat study would not add value**
 229

230 **Rationale**

231 The WoE analysis supports the conclusion that the molecule was sufficiently studied at high

232 exposure margins, and cause-for-concern was not identified for any of the WoE factors.

233

234 **2-year Rat Study Results: No test article related neoplastic findings were present in the 2-year rat**
235 **study.**

236

237 **WoE Criteria**

238

239 Knowledge of intended drug target and pathway pharmacology relative to carcinogenesis

240 • Non-mammalian target excludes intentional alteration of potential mammalian
241 carcinogenic pathways.

242 • No evidence of carcinogenic outcome in 2-year rat studies conducted with other
243 compounds with the same non-mammalian pharmacological target

244

245 Secondary Pharmacology Screen

246 • No evidence of off-target interactions at drug concentrations up to 10 µM, including
247 no interaction with estrogen, androgen, glucocorticoid receptors

248

249 General Toxicology from Chronic Rat Study

250 • Chronic (6-month) toxicology study in Wistar rats dosed to saturation of absorption,
251 achieving up to a 31-fold margin to human exposure.

252 • No evidence of human specific major metabolites.

253 • No treatment-related histopathologic findings observed in standard battery of tissues

254

255 General Toxicology from Chronic Non-rodent Study

256 • Chronic administration (9-month) to non-human primates identified bile duct
257 hyperplasia and hepatocellular hypertrophy, with reactive neutrophils and
258 regenerative hyperplasia. A No-Adverse-Effect-Level was identified which provided
259 a 5-fold margin to human exposure.

260 • Further evaluation in rats would not provide useful information, as similar findings
261 were not observed in the chronic rat study.

262

263 Hormonal Perturbation

264 • No treatment-related findings on reproductive organ weights or histopathology

265

266 Genetic Toxicology

267 • No evidence of genotoxic potential based on criteria from ICH S2(R1) Guidance

268

269 Immune Toxicology

270 • No treatment-related changes in clinical pathology or histopathology of immune
271 tissues (e.g., lymphoid organs, spleen, thymus, bone marrow)

272

273 Additional Special Investigations

274 • No data available

275

276

277 **Case 2: A small molecule antagonist of a neuronal G-protein coupled receptor**

278

279 **Prospective WoE Assessment: Unanimously concluded as likely to be carcinogenic in rats but not**
280 **in humans through well recognized mechanisms known to be human irrelevant, such that a 2-**
281 **year rat study would not add value**

282
283
284
285
286
287
288
289
290
291
292
293
294
295
296
297
298
299
300
301
302
303
304
305
306
307
308
309
310
311
312
313
314
315
316
317
318
319
320
321
322
323
324
325
326
327
328
329
330
331

Rationale

The WoE analysis indicates the potential for rodent-specific liver and thyroid neoplasms based on the toxicology observed in the chronic rat study and on tumor outcome with the pharmacological class. Induction of hepatic cytochrome P450 was demonstrated. Evidence of hormonal perturbation is understood from target pharmacology, did not result in changes in reproductive organ weight or histopathology, and occurred at high multiples to human exposure.

2-year Rat Study Results: The 2-year rat study demonstrated hepatocellular hypertrophy but no neoplastic findings.

WoE Criteria

Knowledge of intended drug target and pathway pharmacology relative to carcinogenesis

- Predominate receptor expression in brain with lower expression in some peripheral tissues, similar across species
- Receptor activation increases ACTH release from pituitary secondary to hypothalamic production of adrenocorticotropin-releasing hormone.
- Hypothalamic receptor ligand levels associated with LH surge and gonadotropin release in rats.
- Target knock-out mice showed no findings related to carcinogenicity.
- Long-term studies with other compound with same pharmacological target associated with thyroid follicular cell adenoma/carcinoma in rats, consistent with elevated thyroid stimulating hormone following off-target cytochrome P450 induction.
- Antagonist binding interaction identified for one off-target receptor with Ki 8-fold higher than Cmax at maximum clinical dose. Known target pharmacology of off-target receptor not associated with tumorigenesis.

General Toxicology from Chronic Rat Study

- Increased liver hypertrophy and organ weight at 50x to 74x margin to human exposure.
- Increased thyroid follicular hypertrophy at 170x to 670x margin to human exposure.
- No evidence of human specific metabolites.
- An active major human metabolite in humans was also present in rats

General Toxicology from Chronic Non-rodent Study

- Increased liver hypertrophy and organ weight at ~230-fold human exposure.

Hormonal Perturbation

- Reduced adrenal weight without histopathological correlates and reduced ACTH level at >74x human exposure in the chronic rat study, consistent with inhibition of drug target. Response noted to be growth suppressive.
- Irregular estrous cycles and decreased pregnancy rate were observed at 60-fold human exposure, and decreased numbers of corpora lutea, implantations, and live embryos were observed at >500-fold human exposure in a fertility study in rats. Considered consistent with inhibition of drug target.
- No treatment-related changes observed in reproductive organ weight or histopathology in chronic rat study.

332 Genetic Toxicology

- 333 • No evidence of genotoxic potential of parent or major human metabolite based on
334 criteria from ICH S2(R1) Guidance

335
336 Immune Toxicology

- 337 • No treatment-related changes in clinical pathology, lymphocyte subsets, or
338 histopathology of immune tissues (e.g., lymphoid organs, spleen, thymus, bone
339 marrow)

340
341 Additional Special Investigations

- 342 • Increased induction of CYP1A2 and CYP3A1 demonstrated
343 • Bone and teeth fluorosis related to defluorination of compound, demonstrated not to
344 occur in humans

345
346
347 **Case 3: A first-in-class small molecule inhibitor of a ubiquitously expressed**
348 **serine/threonine kinase**

349
350 **Prospective WoE Assessment: Unanimously concluded to be uncertain with respect to the**
351 **carcinogenic potential for humans, and a 2-year rat carcinogenicity study is likely to add value to**
352 **human carcinogenicity assessment**

353
354 **Rationale**

355 Significant carcinogenic uncertainty is based on a complex target pharmacology, the lack of
356 precedent with the drug target, and histopathological changes of concern with inadequate
357 mechanistic explanation from the chronic rat study which are supported by similar findings in
358 cynomolgus monkeys. The immune toxicology observed in monkey will contribute to the
359 overall assessment of risk but is not expected to be further informed by a rat carcinogenicity
360 study.

361
362 **2-year Rat Study Results: The 2-year rat study demonstrated an increased incidence, lethality,**
363 **and reduced latency of pituitary tumors in both sexes. This carcinogenic outcome in rats would**
364 **contribute to the overall assessment of human carcinogenic potential.**

365
366 **WoE Criteria**

367
368 Knowledge of intended drug target and pathway pharmacology relative to carcinogenesis

- 369 • Target activation by inflammation-related oxidative stress promotes cellular apoptosis
370 and is linked to control of cell proliferation; target inhibition suppresses apoptotic
371 signaling and impacts cell proliferation, theoretically promoting cancer growth.
372 • Drug target displays tissue-dependent roles in cancer development, both promotion
373 and suppression, in animal models.
374 • No data available on tumor outcome from target inhibition in long term rodent or
375 short term transgenic mouse studies

376
377 General Toxicology from Chronic Rat Study

- 378 • Increased incidence and severity of renal basophilic tubules, eosinophilic droplets,
379 and brown pigment in renal cortex starting at 14-fold human exposure. Etiology of
380 lesions not empirically addressed.

- 381 • Chronic irritation of limiting ridge in non-glandular stomach at 39-fold human
382 exposure. Etiology of lesions not empirically addressed.
- 383 • Increased liver weight without microscopic correlates.
- 384 • No evidence of human specific metabolites.
- 385 • An inactive major human metabolite in humans was also present in rats

386

387 General Toxicology from Chronic Non-rodent Study

- 388 • In monkeys, gastrointestinal epithelial degeneration, necrosis, reactive hyperplasia,
389 ectasia, inflammation, and ulceration, at doses ~12-fold human exposure
- 390 • Increased incidence of renal tubule degeneration /regeneration, necrosis, dilation, and
391 vacuolation at ~12-fold human exposure

392

393 Hormonal Perturbation

- 394 • Increased adrenal weight and cortical hypertrophy in rats at 17-fold human exposure.
395 Etiology not empirically addressed.

396

397 Immune Toxicology

- 398 • In monkeys, suppression of TDAR with no effect on NK cytotoxicity or granulocyte
399 function, and decreased lymphoid cellularity in spleen, thymus, lymph nodes at 12-
400 fold human exposure.

401

402 Genetic Toxicology

- 403 • No evidence of genotoxic potential of parent or major human metabolite based on
404 criteria from ICH S2(R1) Guidance

405

406 Additional Special investigations

- 407 • Increases in hepatic enzymes CYPs 1A, 3A, and 2B demonstrated.

408

409

410 **Case 4: A first-in-class small molecule inhibitor of a prostaglandin receptor**

411

412 **Prospective WoE Assessment: Unanimously concluded as likely not to be carcinogenic in both rats**
413 **or humans such that a 2-year rat study would not add value**

414

415 **Rationale**

416 When compared with the test agent discussed in Case 3, which is also first-in-class, the drug
417 target in Case 4 is not associated with a role in cancer development, histopathological findings
418 were not observed in the chronic rat study, and a large margin of exposure was calculated at
419 the high dose (>50x). The secondary pharmacology screen also indicated the test agent
420 demonstrates target selectivity.

421

422 **2-year Rat Study Results: The 2-year rat carcinogenicity study did not demonstrate a dose-related**
423 **increase in tumors.**

424

425 **WoE Criteria**

426

427 Knowledge of intended drug target biology and pharmacologic mechanism relative to
428 carcinogenesis

ICH S1B(R1) Guideline

- 429
- 430
- 431
- 432
- 433
- 434
- 435
- 436
- Receptor activation associated with allergic inflammatory response and currently available data do not suggest a role in tumor initiation or progression.
 - Knock-out mice of drug target showed no histological abnormalities or effects on immune function during one year of observation.
 - No data available on tumor outcome in 2-year rat studies conducted with other compounds with the same pharmacological target.
 - No data available from a rasH2-Tg carcinogenicity study conducted with the test agent.

437

438 Secondary pharmacology screen

- 439
- 440
- 441
- 442
- 443
- 444
- Test agent was at least 300-fold more selective for drug target when compared with other receptors in the same class as well as a sub-set of other assessed receptors involved in the inflammatory response.
 - Test agent was at least 2000-fold more selective for the drug target in a secondary pharmacology screen of various receptors, ion channels, transporters and enzymes.

445 General Toxicology from Chronic Rat Study

- 446
- 447
- 448
- 449
- Histopathological assessments conducted as part of repeated-dose toxicity studies up to 26-weeks indicated no proliferative changes in any organ or tissue at the highest dose tested (~ 54-fold human exposure based on AUC).
 - No evidence of human specific metabolites.

450

451 General Toxicology from Chronic Non-rodent Study

- 452
- 453
- 454
- Histopathological assessments conducted as part of repeated-dose toxicity studies up to 39-weeks indicated no proliferative changes in any organ or tissue at the highest dose tested (~ 45-fold human exposure based on AUC).

455

456 Hormonal Perturbation

- 457
- No treatment-related findings on reproductive organ weights or histopathology.

458

459 Genetic Toxicology

- 460
- No evidence of genotoxic potential based on criteria from ICH S2(R1) Guidance.

461

462 Immune Toxicology

- 463
- 464
- 465
- 466
- In the 26-week rat toxicity study, there were no effects on immune function (including the TDAR assay evaluating primary and secondary antibody responses) or adverse effects on lymphocyte subsets at the highest dose tested (~54-fold human exposure based on AUC).

467

468 Additional Special Investigations

- 469
- Not performed.